

New Record of *Lipoptena cervi* and Updated Checklist of the Louse Flies (Diptera: Hippoboscidae) of the Republic of Korea

HEUNG CHUL KIM,¹ SUNG TAE CHONG,¹ JOON-SEOK CHAE,² HANG LEE,³ TERRY A. KLEIN,⁴
SANG JAE SUH,⁵ AND LEOPOLDO M. RUEDA⁶

J. Med. Entomol. 47(6): 1227–1230 (2010); DOI: 10.1603/ME09262

ABSTRACT This is the first confirmed record of the genus *Lipoptena* Nitzsch and its species, *Lipoptena cervi* (Linnaeus), from the Republic of Korea. A total of five females and 10 males was collected from eight of 29 Korean water deer, *Hydropotes inermis argyropus* Swinhoe, from Gangwon and Gyeongsangbuk Provinces and Ulsan Metropolitan Area from May through October 2008. An updated checklist of Korean hippoboscids contains nine species in six genera (*Hippobosca* Linnaeus, *Icosta* Speiser, *Lipoptena*, *Ornithoctona* Speiser, *Ornithomya* Latreille, and *Ornithophila* Rondani). Hosts, collection records, and repositories are also noted.

KEY WORDS louse fly, *Lipoptena cervi*, Hippoboscidae, water deer, Korea

Both sexes of louse flies or hippoboscid flies are obligate blood-feeding ectoparasites of birds and mammals. The adults are dorsoventrally flattened and adapted for moving through the bird's plumage and the mammal's pelage, and have specialized claws that aid them to cling to the feathers or hair of their hosts. Worldwide, the family Hippoboscidae consists of three subfamilies, 21 genera, and 213 species (Dick 2006). The genus *Lipoptena* is comprised of ≈30 species (Dick 2006). Members of the genus *Lipoptena*, commonly called deer keds, are winged when they emerge from their puparia, which often remain attached to the pelage of the host or fall to the ground. After emergence, the winged louse flies seek a suitable host, whereby they break their wings off along a cleavage line and become permanently associated with that host. *Lipoptena cervi* has been reported from temperate areas of Europe, Siberia, Northern China, and North America (Maa 1969c). They are obligate parasites of elk, deer, and other bovines, sucking blood

and ovipositing individual prepuparium on the host for up to 10 mo. Although little is known about the vector potential of hippoboscid flies, *L. cervi* are known to harbor and transmit *Bartonella schoenbuchensis* Dehio in deer in Germany and the United States, and animals infested with hippoboscid flies may develop a moderate to severe dermatitis (Dehio et al. 2004, Hermsilla et al. 2006, Matsumoto et al. 2008).

Except for a few early surveys of louse flies from 1922 to 1969 in the Republic of Korea (ROK), no other louse fly surveys of wild birds and large mammals have been conducted in the ROK (Okamoto 1924; Bequaert 1941; Maa 1967, 1969a, b, c; Seok 1970) (Table 1). Only eight species of louse flies were recorded earlier from birds and mammals from the ROK (Maa 1967, 1969c; Anonymous 1994).

In this study, we report the first record of the deer ked, *L. cervi*, collected from the Korean water deer, *Hydropotes inermis argyropus* Swinhoe. We also include an updated checklist of louse flies and their associated hosts in the ROK.

Materials and Methods

Ectoparasites were collected from May through October 2008 from wild deer captured for necropsy from two metropolitan areas, namely the following: Seoul (37.5898 N, 127.0041 E) and Ulsan (Yeampo-dong, 35.5276 N, 129.2924 E; Onyang-eup, 129.2961 E, 35.4234 N) and four provinces, namely the following: Gangwon-do (Cheorwon-gun, 37.0553 N, 127.2139 E), Chungcheongbuk-do (36.8931 N, 127.8469 E), Jeollanam-do (35.1176 N, 126.9367 E), and Gyeongsangbuk-do (Uljin, 37.366009 N, 129.2628 E) in the ROK (Fig. 1), under an approved Korean wildlife animal use protocol, College of Veterinary Medicine, Seoul National University. Louse flies and ticks collected from the deer from above local-

The views expressed in this article are those of the authors and do not reflect the official policy or position of the Department of Defense, the Department of the Army, or the United States Government.

¹ 5th Medical Detachment, 168th Multifunctional Medical Battalion, 65th Medical Brigade, Unit 15247, APO AP 96205-5247.

² Research Institute for Veterinary Science and BK21 Program for Veterinary Science, Veterinary Internal Medicine, College of Veterinary Medicine, Seoul National University, Seoul 151-742, Republic of Korea.

³ Conservation Genome Resource Bank for Korean Wildlife, College of Veterinary Medicine, Seoul National University, Seoul 151-742, Republic of Korea.

⁴ Force Health Protection and Preventive Medicine, 65th Medical Brigade/USAMEDDAC-Korea, Unit 15281, APO AP 96205-5281.

⁵ Department of Environmental Horticulture, College of Life Science and Natural Resources, Kyungpook National University, Sangju 742-711, Republic of Korea.

⁶ Corresponding author: Walter Reed Biosystematics Unit, Division of Entomology, Walter Reed Army Institute of Research, 503 Robert Grant Avenue, Silver Spring, MD 20910-7500.

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 2010		2. REPORT TYPE		3. DATES COVERED 00-00-2010 to 00-00-2010	
4. TITLE AND SUBTITLE New Record of Lipoptena cervi and Updated Checklist of the Louse Flies (Diptera: Hippoboscidae) of the Republic of Korea			5a. CONTRACT NUMBER		
			5b. GRANT NUMBER		
			5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S)			5d. PROJECT NUMBER		
			5e. TASK NUMBER		
			5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Walter Reed Army Institute of Research, Walter Reed Biosystematics Unit, 503 Robert Grant Avenue, Silver Spring, MD, 20910			8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSOR/MONITOR'S ACRONYM(S)		
			11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT This is the 1st confirmed record of the genus Lipoptena Nitzsch and its species Lipoptena cervi (Linnaeus), from the Republic of Korea. A total of 9 females and 10 males was collected from eight of 29 Korean water deer, Hydropotes inermis argyropus Swinhoe, from Gangwon and Gyeongsangbuk Provinces and Ulsan Metropolitan Area from May through October 2008. An updated checklist of Korean hippoboscids contains nine species in six genera (Hippobosca Linnaeus, Icosta Speiser, Lipoptena, Ornithoctona Speiser, Ornithomya Latreille, and Ornithophila Rondani). Hosts, collection records, and repositories are also noted.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 4	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Table 1. Checklist of the louse fly (Hippoboscidae, Diptera) species and associated hosts from the Republic of Korea

Species	n	Host	Type	Collection date	Location	Repository (code)	References
<i>Hippobosca equina</i> Linnaeus	NR	NR	Mammal	NR	Jeju Island, Jeju-do	NR	Seok (1970)
<i>H. longipennis</i> Fabricius	1 F	<i>Motacilla alba leucopsis</i> Gould	Bird	1925	Taiyudong	HKU	Maa (1967)
	1 F	<i>M. alba leucopsis</i>	Bird	May 1922	Seoul	HKU	Maa (1967)
	6	<i>M. alba leucopsis</i>	Bird	June 1922	Quelpart Island (=Jeju), Jeju-do	HKU	Okamoto (1924)
<i>Icosta ardae ardae</i> (Macquart)	1 M	<i>M. alba leucopsis</i>	Bird	NR	Chemulpo (=Inchon), Gyeonggi-do	BMNH	Maa (1969b)
<i>Lipoptena cervi</i> (Linnaeus)	1 F, 2 M	<i>Hydropotes inermis argyropus</i> Swinhoe	Mammal	8 May and 16 June 2008	Cheorwon-gun, Gangwon-do	FMDL	X
	3 M	<i>H. inermis argyropus</i>	Mammal	16 June 2008	Ulsin-gun, Gyeongsangbuk-do	FMDL	X
	1 F, 1 M	<i>H. inermis argyropus</i>	Mammal	5 Aug. and 10 Oct. 2008	Yeampo-dong, Buk-gu, Ulsan	FMDL	X
<i>Ornithoctona plicata</i> von Olfers	3 F, 4 M	<i>H. inermis argyropus</i>	Mammal	21, 23 Aug. 2008	Onyang-eup, Ulsu-gun, Ulsan	FMDL, SI (1 F)	X
	NR	NR	NR	NR	Unknown location, Korea	NR	Bequaert (1941), Maa (1969c)
<i>Ornithomyia acicularia aobatonis</i> Matsumura	2 F	<i>Emberiza spodocephala</i> Pallas	Bird	Oct. 1965	Kunja	BBM (75079)	Maa (1967)
	1 F	<i>E. spodocephala</i>	Bird	Oct. 1965	Chin Chup Myun, Kyunggi-do	MAPS (6E 1044)	Maa (1967)
	1 F	<i>E. tristrami</i> Swinhoe	Bird	Oct. 1965	Nao Chon Myun, Kyunggi-do	MAPS (6E 0851)	Maa (1967)
	1 F	<i>E. rutila</i> Pallas	Bird	Oct. 1965	Chin Chup Myun, Kyunggi-do	MAPS (6E 1370)	Maa (1967)
	1 F	<i>Acrocephalus arundinaceus</i> (Temminck and Schlegel)	Bird	Aug. 1965	Chin Chup Myun, Kyunggi-do	MAPS (6E 1004)	Maa (1967)
<i>O. chloropus extensa</i> Maa	1 F	<i>E. rutila</i>	Bird	Oct. 1966	Chin Chup Myun, Kyunggi-do	MAPS (6E 1467)	Maa (1967)
<i>O. fringillina</i> Curtis	1 F	<i>Coccothraustes coccothraustes japonicus</i> Temminck and Schlegel	Bird	Nov. 1965	Seoul	NR	Maa (1967, 1969c)
<i>Ornithophilla metallica</i> (Schiner)	1 F	<i>M. alba leucopsis</i>	Bird	NR	Seoul	NR	Maa (1969a)

BBM, Bernice Bishop Museum, Honolulu, HI; BMNH, British Museum of Natural History, London, United Kingdom; F, female; FMDL, 5th Medical Detachment Laboratory, United States Army, Seoul, ROK; HKU, Entomology Institute of Hokkaido University, Sapporo, Japan; M, male; MAPS, Migratory Animal Pathology Survey, Honolulu, HI; NR, no available record; SI, National Museum of Natural History, Smithsonian Institution, Washington D.C.; X, this survey.




Fig. 1. Map of the ROK showing collection localities of the louse fly, *L. cervi* (triangles, with collected flies; circles, without flies; *n* = number of louse flies/number of infested hosts/number of examined hosts), collected from the Korean water deer, *H. inermis argyropus*. In Jeju Island, only roe deer, *C. pygargus*, was captured. (1) Cheorwon-gun, Gangwon-do (*n* = 3:2:4); (2) Uljin-gun, Gyeongsangbuk-do (*n* = 3:1:2); (3) Yeampo-dong, Buk-gu, Ulsan (*n* = 2:2:2); (4) Onyang-eup, Ulju-gun, Ulsan (*n* = 7:3:9); (5) Seoul (*n* = 0:0:1); (6) Chungcheongbuk-do (*n* = 0:0:2); (7) Jeollanam-do (*n* = 0:0:7); (8) Jeju Island, Jeju-do (*n* = 0:0:1).

ities were placed in 2-ml plastic vials with 80% ethyl alcohol and transported to the 5th Medical Detachment laboratory, where louse flies were identified under a dissecting microscope using morphological keys (Bequaert 1941; Maa 1965, 1967). Specimens were sent to the Department of Entomology, Smithsonian Institution, where the identifications were confirmed by Dr. F. C. Thompson. A voucher specimen (one adult female) was deposited at the National Museum of Natural History, Smithsonian Institution (Washington D.C.), whereas the rest of the identified samples were returned to Seoul National University for assay of selected zoonotic pathogens.

Results and Discussion

A total of 29 Korean water deer (*H. inermis argyropus*) and one roe deer (*Capreolus pygargus* Pallas) was captured from May through October 2008. A total of 15 adults (five females and 10 males) and two puparia of *L. cervi* was collected from eight Korean water deer at Gangwon and Gyeongsangbuk Provinces, and the Ulsan Metropolitan Area (Table 1; Fig. 1). Previously, only species belonging to five genera (*Ornithoctona*, *Ornithomya*, *Ornithophilla*, *Icosta*, and *Hippobosca*) of the family Hippoboscidae were reported from the ROK (Table 1).

In addition to the first record of *L. cervi* from three widely dispersed geographical locations extending from the northern demilitarized zone to the southernmost part of the ROK, we also report for the first time the occurrence of the genus *Lipoptena* in this country. Adult *L. cervi* is dorsoventrally flattened, relatively small (5–7 mm long), and brownish in color (Fig. 2). The head has distinct simple eyes (or ocelli) and well-developed maxillary palps. The thorax has 15–18 laterocentral setae on the mesonotum with spines scattered on the anterior part of prosternal lobe. Both sexes alight on a host, take a blood meal, mate, and after an indeterminate period the female will give birth to mature larva that will begin to pupate (<http://ento.psu.edu/extension/factsheets/deer-keds>). In the present survey, two puparia were collected while they were attached to the pelage of deer, and the wings of all adults were broken off close to their bases.

The changing ecology (particularly reforestation) and increases in associated wildlife populations provide for great proximal host contacts and wide distribution of blood-feeding ectoparasites in the ROK. Further studies are needed to identify the geographical distributions of louse flies, including *L. cervi*, and their potential role as reservoirs of the zoonotic pathogens in the ROK. With this finding of the new occur-




Fig. 2. Adult male *L. cervi* collected from the Korean water deer (*H. inermis argyropus*) in the ROK. (A) Dorsal view; (B) ventral view.

rence of *L. cervi* from the two provinces (Gangwon-do, Gyeongsangbuk-do) and a metropolitan city (Ulsan), prediction models may be developed to determine the potential distribution of this louse fly species, its associated mammalian host, and the risk assessment of possible hippoboscids-borne human and animal diseases in the Korean peninsula.

Acknowledgments

We thank Captain Peter V. Nunn, 5th Medical Detachment, 168th Medical Battalion, 65th Medical Brigade, and the Conservation Genome Resource Bank for Korean Wildlife, Seoul National University, for their support in this work. We also thank Jun-Gu Kang and Sungjin Ko, College of Veterinary Medicine, Seoul National University, for collection of deer keds in this survey; and Richard Robbins, Armed Forces Pest Management Board, and Joel Gaydos, Armed Forces Health Surveillance Center for their support in conducting this work. Special thanks go to F. C. Thompson for confirming the identification of the specimens, and to D. M. Levin and B. P. Rueda for reviewing the manuscript and providing helpful suggestions. Portions of this work were supported by National Research Foundation of Korea grant funded by the Korean Government (2009-0071622), the Armed Forces Health Surveillance Center, Global Emerging Infections Surveillance and Response Systems (Silver Spring, MD), and the National Center for Military Intelligence (Fort Detrick, MD).

References Cited

- Anonymous. 1994. Check list of insects from Korea. Konkuk University Press, Seoul, Korea.
- Bequaert, J. 1941. The Hippoboscidae of Oceania (Diptera). Occas. Pap. Bernice P. Bishop Museum, Honolulu, Hawaii 16: 247–292.
- Dehio, C., U. Sauder, and R. Hiestand. 2004. Isolation of *Bartonella schoenbuchensis* from *Lipoptena cervi*, a blood-sucking arthropod causing deer ked dermatitis. J. Clin. Microbiol. 42: 5320–5323.
- Dick, C. W. 2006. Checklist of world Hippoboscidae (Diptera: Hippoboscoidea). Department of Zoology, Field Museum of Natural History, Chicago, IL.
- Hermosilla, C., N. Pantechev, R. Bachmann, and C. Bauer. 2006. *Lipoptena cervi* (deer ked) in two naturally infested dogs. Vet. Rec. 159: 286–287.
- Maa, T. C. 1965. A synopsis of the Lipopteninae (Diptera: Hippoboscidae). J. Med. Entomol. 2: 233–248.
- Maa, T. C. 1967. A synopsis of Diptera pupipara of Japan. Pac. Insects 9: 727–760.
- Maa, T. C. 1969a. Synopses of the genera *Ornithophila* and *Ornithoctona* with remarks on their habitat diversification (Diptera: Hippoboscidae). Pac. Insects Monogr. 20: 1–23.
- Maa, T. C. 1969b. Revision of *Icosta* (= *Lynchia* Auctt.) with erection of a related genus *Phthona* (Diptera: Hippoboscidae). Pac. Insects Monogr. 20: 25–203.
- Maa, T. C. 1969c. A revised checklist and concise host index of Hippoboscidae (Diptera). Pac. Insects Monogr. 20: 261–299.
- Matsumoto, K., Z. L. Berrada, E. Klinger, H. K. Goethert, and S. R. Telford. 2008. Molecular detection of *Bartonella schoenbuchensis* from ectoparasites of deer in Massachusetts. Vector Borne Zoonotic Dis. 8: 549–554.
- Okamoto, H. 1924. The insect fauna of Quelpart Island (Saishiu-to). Bull. Agr. Exp. Stat., Gov. Chosen 1: 47–233.
- Seok, D. M. 1970. The insect fauna of the Island Quelpart. Po-Chin-Chai Co, Seoul, Korea.

Received 2 November 2009; accepted 16 July 2010.